

Asymptotics of the Infrared

P.R. Crompton

*Institut für Theoretische Physik, Universität Leipzig,
Augustusplatz 10/11, D-04109 Leipzig, Germany.**

(Dated: February 1, 2008)

We follow recent formulations of dimensionally reduced loop operators for quantum field theories and exact representations of probabilistic lattice dynamics to identify a new scheme for the evaluation of partition function zeroes, allowing for the explicit analysis of quantum critical phenomena. This new approach gives partition function zeroes from a factored quantum loop operator basis and, as we show, constitutes an effective mapping of the renormalization group β -function onto the noncommuting local operator basis of a countably finite Hilbert space. The Vafa-Witten theorem for CP-violation and related complex action problems of Euclidean Field theories are discussed, following recent treatments, and are shown to be natural consequences of the analyticity of the limiting distribution of these zeroes, and properties of vacuum regimes governed by a dominant quantum fluctuation in the vicinity of a renormalization group equation fixed point in the infrared.

PACS numbers: 73.43.Nq, 11.15.Ha, 11.10.Gh.

A classical spin system and its quantum system analogue apparently share many similarities at the level of the spin Hamiltonian. The difference between quantum and classical formulations is perhaps somewhat subtle at first glance. In the latter case the spin components are represented through the action of an operator basis, rather than purely orthogonal vector basis elements as is the case with the former. Although the dimensionality of a particular model is unaffected by this step of operator inclusion the partition function of the quantum system is formally defined as a trace over this operator basis acting on some suitably structured Hilbert space (whereas the classical partition function is strictly given simply through a summation over spin vector realisations). This inclusion therefore adds an extra dimension to evaluation of the quantum partition function that maybe then dealt with, or expressed, through a variety of scenarios relating to the applicability of dimensional reduction. This basis reformulation or inclusion of operator components can in fact be argued to be a form of second quantisation of a given classical model [1].

We are ultimately interested here to derive an exact polynomial expansion for the partition function for a generic quantum system formed in such an operator basis, and to discuss the effect that quantum fluctuations have on frustrating this effort. The zeroes of this polynomial expansion then constitute the partition function zeroes whose limiting distribution we aim to now analyse following the arguments and reasoning presented in [2][3]. However, the difference from these two cases we will argue (and would hope to stress as a central result) is that in general we identify the partition function of a generic quantum system is nonanalytic and unamenable to such

expression. Our reasoning will follow much as for the general case of renormalization group equations of motion and the β -function, which may become singular prior to reaching its asymptotic value through infrared divergences [4]. Crucially dependent on structuring the formal limits in the countably finite Hilbert space basis we will argue a limiting analyticity can be defined, however, using recent results and discussions of the structure of the poles of the inverse Laplace transform in the context of recent results on exact lattice probabilistic dynamics [5][6].

It may initially appear somewhat confused to attempt to define bounds for the analyticity of this expansion, or indeed of little practical relevance to order and arrange a form for an exact expansion that is valid essentially only in the vicinity of the continuum limit. This is certainly somewhat of an anathema to the scaling arguments of the renormalization group and partition function zeroes formulations where scale invariance is a central point. Our major thrust, however, will be to treat disconnected contributions to the partition function in a local basis, marking a certain departure from (quasi-classical) perturbative expansion. The noncommutativity of the operators for the quantum system force the reformulation in this regard. Diagrammatically disconnected contributions will be here suppressed through the limiting distributions of the basis operators formed on the Hilbert space, rather than through the tacit convergence of any effective series. Akin to partition function zeroes formulation, singularities are expressed through the limiting asymptotic behaviour of poles as the continuum is approached, but here with an intermediate result closely connected with dimensional reduction that factorises a local symmetry preservation. To facilitate this discussion we first review salient points in the derivation of the transfer matrix and associated Quantum Monte Carlo method for assigning exact probabilistic dynamics to a loop operator basis, al-

*Electronic address: crompton@itp.uni-leipzig.de

though we would stress this is not primarily intended as a discussion to motivate specific numerical approaches.

I. NONCOMMUTATIVITY OF THE HAMILTONIAN OPERATOR

A number of algorithmic scenarios exist for the Monte Carlo simulation of quantum spin systems [7][8][9][10][11]. They were developed for the purpose of addressing the critical slowing down that affects the simulation of quantum systems at low temperatures. They differ largely from standard lattice gauge theory simulation methods, for example, in that in the main they are driven locally. Probabilistic Monte Carlo decisions are taken on a lattice site-by-site basis rather for the global lattice configuration as is usually the case with the latter [12]. Although, we will later comment on the relevance of the results to the complex action problem and recent lattice attempts at a general solution in this direction here algorithms for the numerical study of quantum spin systems serve as a means to introduce the formulation a complete-orthonormal operator basis acting on a Hilbert space in the context of exact lattice probabilistic dynamics. Following recent discussion on the limiting continuum distribution of a suitably arranged Markov process, we then argue this is amenable to analytic expression. Thus a means is identified to formulate partition function zeroes to express singular behavior in a quantum system for cases where otherwise the non-commutativity of the quantum operator will basis spoils this analyticity. This result is so expressed through an approach closely connected to dimensional reduction [1]. From this derivation it becomes then possible to the define bounds under which a generated Monte Carlo distribution becomes suitably close to the continuum result that the zeroes of the partition function can be meaningfully associated with the limiting thermodynamic arguments for the singularities of [2][3]. Or more properly now their quantum system analogues, as we will argue from the results on the behaviour of the poles of the inverse Laplace transform of this expansion. In this sense, therefore, determining the statistical limitations of numerical Monte Carlo sampled loop operator distributions with relevance to this task remains an onerous and somewhat separate ongoing direction, although we return to discussion of the complex action problem in a later section.

Proceeding with commenting on generic Quantum Monte Carlo approaches for quantum spin systems, the pedagogical example usually taken is the spin one half anti-ferromagnetic Heisenberg model ($S = 1/2$ AFM)[7][9]. The transfer matrix of this system contains the simplest off-diagonal singlet contribution possible to express a quantum interaction. The spins defined $S_i = 1/2 \sigma_i$ are defined for the model on each lattice site index i of a d -dimensional hypercube, with a nearest-neighbour interaction and σ_i being the Pauli matrices. In general, ie. for all but vanishing off-diagonal

contributions of the suitably defined transfer matrix, the spin operators do not commute. If one were to wish to evaluate the Boltzmann factor exactly from the nearest-neighbour spin Hamiltonian thus defined on a finite lattice volume this would then present something of a problem, akin to the difficulties of representing Grassmann algebras in lattice gauge field theories. The combined idea of Suzuki and Trotter as a basis for numerical simulation was to partition the nearest-neighbour Hamiltonian operator $\hat{\mathcal{H}} = \sum_i H_i$. For the case of the $S=1/2$ AFM the leading result would be then given for an odd-even site partitioning of the operators.

$$e^{-\beta \hat{\mathcal{H}}} \approx \prod_i \exp(-\Delta\tau H_i) + \mathcal{O}(\Delta\tau^2) \quad (1)$$

The error in estimating the exponential is thus subdominant despite the noncommutativity of the nearest-neighbour spin operators that constitute the Hamiltonian $\hat{\mathcal{H}}$. This is dependant evidently on the value of $\Delta\tau$, as yet undefined. The partition function itself is defined as the trace over the basis on which the operators act, and thus the extent of the above product i must clearly be somehow dependent on the lattice volume of the classical spin model on which the quantum system operators are introduced. For the $S=1/2$ AFM, as we have said, the leading result is given through odd-even partitioning. There would be no reason a priori, though, to define a particular lattice extent for the additional dimension introduced by the operators themselves. Indeed, only two points need be formally defined for the trace operation. If $\Delta\tau = \beta/L_t$ is the defined as the lattice spacing in Euclidean time the only necessary points to preserve the measure are $\Delta\tau = 0$ and $\Delta\tau = \beta$. However, we wish to discretise the Euclidean time direction, not simply for the purposes of numerical simulation but to now enable discussion of the recent limiting results of probabilistic dynamics on this Euclidean extent by first treating it as a countably finite space. To properly define the partition function $Z(\beta)$, it is necessary to first note that the elements of the factorised product are the transfer matrices that act on the suitably arranged complete-orthonormal set of basis states.

$$Z(\beta) \equiv \text{Tr } e^{-\beta \hat{\mathcal{H}}} \approx \sum_{\{S_{it}\}} \prod_p W_p(\{S_p\}) \quad (2)$$

$$W_p(S_p) = \langle S_{i,t} S_{i+1,t} | e^{-\Delta\tau H_i} | S_{i,t+1} S_{i+1,t+1} \rangle \quad (3)$$

The summation of the weights W_p for the transfer matrices above therefore runs over the spin configuration space of the lattice, as given through the state of the elementary plaquettes p . For the $S=1/2$ AFM these are defined as the two-by-two elements of the hypercube defined over a unit spatial i and Euclidean time t extent. The weight W_p associated with the of each plaquette are given by the matrix elements, and can be built up numerically through the probability density function distributions generated

by a Monte Carlo procedure, as we will later discuss [10]. The definition of the transfer matrix elements and their explicit relation to the noncommutativity of the spin operators will also be postponed to later section, whilst we consider an alternate quasi-classical approach [11][13].

The reverse of the above product expansion is of course possible. Rather than proceeding from a local, and inherently limiting-case factorisation perspective, one can aim to treat the statistical fluctuations of the noncommuting operator basis in a global manner, using the ideas of stochastic field theory. Statistical cumulants $\langle H_1 \dots H_N \rangle^c$ can be defined on the operator basis through the differentiation of some suitable generating function,

$$\langle H_1 \dots H_N \rangle^c = \frac{\partial}{\partial \lambda_1} \dots \frac{\partial}{\partial \lambda_N} \ln \langle e^{\lambda_1 H_1} \dots e^{\lambda_N H_N} \rangle \Big|_{\lambda_1 = \dots = \lambda_N = 0}. \quad (4)$$

It is important to note that no assumption is being made for the distribution of the operators on the space in this. If this were the case one would arrive at a standard time-ordered path integral formalism for quantum field theory generating functionals, albeit on a countably finite space (a Fock space definition), rather than this statistical result which is simply for functions rather than functionals. That the time-ordered result is not guaranteed is somewhat obvious since the operators in the cumulant definition are as yet rather general and undefined. Their action could be highly nonlocal or could indeed be incomplete in some sense, in which case the dynamics would be beset by singularities. That the action of a proper Hamiltonian necessarily yields a time-ordered product is somehow less obvious, but will we again postpone a fuller discussion of issues of locality and noncommutativity to the section that follows, where the role of a Markov process in defining exact probabilistic dynamics for the lattice is made evident.

Following this analogy, though, the expectation value of the product defined without the logarithm in the right-hand side would correspond to analogue of the mixture of disconnected and connected n -point functions one obtains from the action of the generating functional in the field theory analogue. Again, without assuming form for the distributions one can take the continuum limit of the operator space $N \rightarrow \infty$ for the above definition of cumulants. This then yields a quasi-classical effective Hamiltonian $\hat{\mathcal{H}}_c$ from equating the cumulant definition to an exact series expansion for an exponential in this limit. This appears evidently simpler to evaluate, being reduced to a quasi-classical formulation and therefore amenable to a summation over vector realisations.

$$Z = \int \mathbf{n} \langle \mathbf{n} | \exp(-\beta \hat{\mathcal{H}}) | \mathbf{n} \rangle \quad (5)$$

$$\exp[-\beta \hat{\mathcal{H}}_c(\mathbf{n}, \beta)] \equiv \langle \mathbf{n} | \exp(-\beta \hat{\mathcal{H}}) | \mathbf{n} \rangle \quad (6)$$

Caution is in order however as one must now expand in the matrix elements of powers of the full Hamiltonian $\hat{\mathcal{H}}$, for a practical numerical approach to evaluate the quasi-classical Hamiltonian $\hat{\mathcal{H}}_c$.

$$\beta \hat{\mathcal{H}}_c(\mathbf{n}, \beta) = \langle \mathbf{n} | \hat{\mathcal{H}} | \mathbf{n} \rangle^c - \frac{\beta}{2!} \langle \mathbf{n} | \hat{\mathcal{H}}^2 | \mathbf{n} \rangle^c + \frac{\beta^2}{3!} \langle \mathbf{n} | \hat{\mathcal{H}}^3 | \mathbf{n} \rangle^c + \dots \quad (7)$$

As was case with the Trotter-Suzuki approach the non-commutativity of the basis cannot be entirely circumvented. This expansion may appear somewhat straight forward, but $\hat{\mathcal{H}}_c$ itself is strictly defined only through the cumulants (connected components), rather than the statistical expectations (connected and disconnected components). A numerical evaluation of the quasi-classical system following similar truncation approximations to the Trotter-Suzuki form therefore suffers from three sources of systematic error. The expansion itself may diverge through either the value of β or the level-spacing diverging as a function of the expansion, and the stochastic expansion terms themselves contain the statistical analogue of disconnected contributions to the quasi-classical partition function in all but the continuum limit. Without unduly pessimistic focus on the numerical prospects, both of the two parallel approaches fall on this issue of noncommutativity. Yet at the same time, this comparison as the continuum limits are separately approached is revealing.

To make a further association with issues of dimensional reduction in gauge field theories we can aim to relate quasi-classical statistical and functional terminologies for effective actions. At small inverse temperature β , the interaction coupling of gauge fields becomes small through asymptotic freedom and perturbative expansion is justified [14][15]. For length scales much larger than those of the Euclidean time extent it also seems reasonable to integrate out the nonstatic (time-dependent) modes of the gauge fields. This gives the leading behavior at small β , but it is important that the perturbative expansion of this dimensionally reduced model does not have divergences in the infrared. The nonstatic modes are suppressed and renormalize the expansion terms of the new effective theory. To develop the above statistical association, the expansion is again valid if the perturbative coupling is not so large as to cause divergence. By way of contrast no explicit assumption is made about the locality of the expansion in the statistical case and so it is not renormalized. The dimensional reduction of the field theory here forms the functional analogue of the stochastic approximation. The functional approach becomes again invalid if the contribution of higher order n -point functions (either connected or disconnected) becomes dominant.

It is essentially the locality of the expansion of the effective action that crucially determines the response to

infrared divergences and so we seek a different approach as a basis for study. By following the Trotter-Suzuki decomposition, as opposed to the effective quasi-classical Hamiltonian formalism, we work in a nonlocal framework without implicit effective series truncation. Lee and Yang, in the context of classical spin systems defined a limiting analyticity for the partition function such that thermodynamic singularities were only strictly expressed through the limiting distributions of the zeroes of polynomials so constructed [2]. We will argue that this specific analyticity property of the partition function is a nontrivial consequence of the renormalization group approach in the quantum case by working over this extended phase space. By addressing the form of the limiting distribution taken by noncommuting operators on the basis space we will seek to similarly define a limiting analyticity for the infrared divergences connected with the renormalization group equation fixed points of quantum systems through this factorisation approach.

II. LOOP-CLUSTER OPERATORS

Before proceeding to discussion of the partition function polynomial form for the quantum case we continue first outlining the relationship between the factorised Hamiltonian operators and their distributions on a Hilbert space. This is the relationship defined through the transfer matrix elements of these operators. We motivate these now necessary definitions through part-review of numerical Quantum Monte Carlo loop-cluster methods. Numerical motivation perhaps seems an unnecessarily heuristic path to take here for the proposing of an exact continuum form for the generic partition function of a quantum system. To reiterate, indeed the stated aim is not to define numerical grounds for simulation. By constructing, or reviewing, the Markov process definition we will simply articulate the complete-orthonormal nature of the basis formed by the elementary plaquettes defined over the spatial and temporal sites, and exact probabilistic dynamics defined thereon.

Again whilst a factorisation property is clearly a key element in properly defining the zeroes of a polynomial, the direct connection with a complete-orthonormal basis for a Hilbert space and partition function zeroes of a generic quantum system would perhaps appear obtuse at this point. However, mirroring Lee and Yang's discussion we associate the analyticity of an exact expansion with identifying the continuous distribution or functional assignment of solutions to the generating functions of said expansion. That each expansion coefficient in the polynomial expansion can represent a well-defined integral equation is actually a necessary starting point of Lee and Yang's discussion [2]. We therefore do this explicitly here by considering the action of taking the vanishing Euclidean time lattice spacing limit for the loop operators on a finite spatial lattice extent. The construction of the quantum system analogue of Lee and Yang's integral

assignment is then given through this action, implicit to the loop-cluster method definition.

The loop-cluster method for quantum systems is similar in form to that of the Swendsen-Wang cluster algorithm of classical statistical mechanics [17][7][9]. The partition function is defined over a set of spin configurations $\{\mathcal{S}\}$ of the lattice, and a set of graphs $\{G\}$ linking these spin states, defining an extended phase space. For the Monte Carlo process statistical weight functions $W(\mathcal{S}, G)$ must be assigned for each spin configuration of a lattice ensemble, such that now all elements of the extended phase space are defined as either being connected or disconnected. Given some configuration of the spins $\mathcal{S}$ of a lattice each bond linking spins is separately considered and assigned such a status based on the value of $W(\mathcal{S}, G)$, determined through the local spin configuration of $\mathcal{S}$. This then implicitly defines a path, or graph subset G , through the volume of the particular configuration. The total weight of a given configuration $W(\mathcal{S})$ in the lattice ensemble is given by the sum over graph weights.

$$\sum_G W(\mathcal{S}, G) = W(\mathcal{S}) \quad p(\mathcal{S} \rightarrow (\mathcal{S}, G)) = \frac{W(\mathcal{S}, G)}{W(\mathcal{S})}, \quad (8)$$

where $p(\mathcal{S} \rightarrow (\mathcal{S}, G))$ is probability of a obtaining a particular graph within a lattice configuration. This then implies both that $W(\mathcal{S}) \neq 0$ and $W(\mathcal{S}, G) \neq 0$. The process is otherwise nonergodic and there is some absolute preferred configuration state of the volume that once reached cannot be left in an exact statistical sense. We are interested to treat singularities, of course, but do not anticipate divergences such as these on a finite lattice for essentially geometric reasons - without, though, wishing to necessarily restrict the spaces of interest at present. The loop-cluster algorithm itself is clearly of more general construction than its applicability to quantum systems, but the Hamiltonian operation we are interested in, acting locally on suitably defined plaquettes, can satisfy this latter Monte Carlo probabilities detailed balance condition.

Returning to the importance of locality. Considering the pedagogical example of the $S = 1/2$ AFM according to the above definition of a graph a single spin cannot be inverted alone on an elementary plaquette to create a valid change between successive lattice configurations. This follows from the implicit local conservation of S . The graphs so defined thus map relative change across the extended volume. In fact, demanding on a given a graph that the form of $W^{plaq}(\mathcal{S}, G)$ does not change at all between consecutive lattice configurations in an ensemble then implies that all allowed plaquette changes match up between neighbouring plaquettes across the volume.

An extensive literature exists on the definition of suitable definition of transfer matrices for quantum systems and the loop-cluster method, but to clarify further for the

pedagogical example the here plaquette has eight spin configurations, $S_p = i^\pm$, of the four site vertices, i [7]. Each transition $i \leftrightarrow j$ defines a respective transfer matrix element $w^{ij} \equiv w^{ji}$, and corresponds in this case to inverting either two or all four of the spins of a plaquette. We can then therefore think to write the partition function for this, or a generic quantum system, in terms of an equivalent bond or loop operator h_b basis defined only in terms of the pertinent transfer matrix elements [7][16][18]. The pertinent transfer matrix elements will be those are nonzero. The diagonal elements of these loop operators for the pedagogical case will correspond to four inversions and comprise the identity operator, and the self-adjoint off-diagonal remainder will be given uniquely through the graphs determined for each configuration (or rather their ensemble-average). This all then essentially represents an effort to define the pseudo-reduction step on a finite basis, mapping the discrete distribution of graphs to the limit of vanishing Euclidean time lattice spacing where a continuous distribution is recovered. The partition function in terms of the loop operators $\hat{h}_b$ is,

$$\text{Tr } e^{-\beta \hat{\mathcal{H}}} = e^{\beta \sum_b J} \lim_{\Delta t \rightarrow 0} \left(\prod_b e^{(-J + J h_b) \Delta t} \right)^{\beta / \Delta t} \quad (9)$$

$$= e^{\beta \sum_b J} \int \rho(d\omega) \prod_b^* h_b, \quad (10)$$

where the Hamiltonian is rewritten over the bonds b as $\hat{\mathcal{H}} = -\sum_b J h_b$ and J is the nearest neighbour coupling. Here the $\prod_b^* h_b$ is a time ordered product of the loop operators with $\rho(d\omega)$ a Poissonian probability measure of the connection status recovered by taking the limit of vanishing Euclidean time lattice spacing. That surprisingly this relatively general construction of the loop operator transfer matrices yields a precise continuous distribution in the limiting step follows directly from the locality property. It is important to note as well that no noncommutativity property is invoked to obtain the Poissonian distribution on the basis elements of the now countably finite spatial element space. One could have equally well ignore the off-diagonal elements altogether and used the approach to map a classical system into another classical system, since after all the method is classical in origin.

The advantage over the quasi-classical series expansion approach here is that the reduced factorised basis elements are explicitly highly nonlocal in character whilst locally preserving S exactly. The central assumption, however, is again that of the quasi-classical expansion: the statistical averages only become interchangeable with those of the thermodynamic or dynamic averages in the infinite volume limit. Here the definition is somehow less ad hoc and the treatment of disconnected contributions less tacit, since locality is enforced site-by-site, which allows us define a polynomial with a slightly different relation to the poles of the partition function. By construction, the basis of plaquettes is complete, and orthonormal and more importantly the weight functions associated to graphs are unique and independent of the

location within the Markov chain or statistical ensemble. Although S is locally preserved the elements of the extended space reached by a finite Markov process can be necessarily incomplete and so functional solution is not necessarily admitted. We could be trite and label this the ambiguities of numerical determination, but the issue of whether or not a discrete extended basis can form a covering of the target space is a perhaps a more interesting question which we will return to shortly.

The extent to which the above partition function corresponds with the Nambu definition of a generalised Hamiltonian (in the operation of elements on a Poisson manifold [19]), and could perhaps then therefore be now further generalised for non-Euclidean metrics, is certainly an interesting question but not one we will aim to further address here. We wish to now investigate the limiting analyticity of this factorised reduced expansion, noticing both that the probability distributions of the operators become exact and continuous in the limit of vanishing Euclidean time lattice spacing and that a product of two-by-two self-adjoint operators (for the $S = 1/2$ AFM case, and of generic form) is particularly simple to diagonalise. We have now identified the continuous distributions associated with factored product which we are aiming to now diagonalise, in keeping with Lee and Yang's motivations, as a first step towards establishing the analyticity properties of the eigenvalues and characteristic polynomial associated with this operator product.

III. PARTITION FUNCTION ZEROES POLYNOMIAL

We have stressed, perhaps seemingly unnecessarily, that the partition function zeroes for a generic quantum system are somehow different than those introduced by Lee, Yang and Fisher which we interpret now as classical results. By considering the action of loop operator elements on an extended phase space, defined through a complete-orthonormal local basis, we have discussed in the previous section the identification of an exact factorised operator product on a countably finite Hilbert space with an associated continuous time-ordered distribution structuring. This is a limiting result, but as we have noted the finite basis loop operators by construction are trivially diagonalisable in general. The new step we now take is to note that there is a necessarily a difference between diagonalising the elements of a direct product and determining the characteristic equation of that product. Herein, we believe, lies the divide in the arguments concerning deriving an exact analytic form for the partition functions of the respective quantum and classical systems.

If the distribution obtained through projecting the graph results onto the spatial extent of the lattice is discrete, rather than continuous, in the quantum case some of the loop operator elements may be defined only on a subspace of the extended space. The transfer matrix

allows for the lattice to be entirely comprised of disconnected contributions, of course, but this singularity is under defined in the discrete quantum case. The eigenvalues of the factored product are therefore not necessarily analytically connected on the finite Hilbert space defined over the spatial elements with a discrete statistical distribution, ie. at finite Euclidean time lattice spacing. Conversely with what we here deem the classical result (if this pictured simply as a limiting case of commuting operators) all elements are necessarily connected through the diagonality of all the terms of the factored product. There may be still disconnected contributions to the partition function but the space of the partition function itself is not now under defined as a consequence, since the plaquettes are constructed by definition to preserve a local symmetry. Up to singularities in the continuum limit for the Hilbert space, as defined over the loop operator basis of the lattice spatial extent, we can thus be assured that the classical case is always defined whereas in the general quantum case is not. At least, to be clear, if we are now aiming to treat the continuum limit of the eigenvalues of our product as the polynomial zeroes of the partition function of Lee, Yang and Fisher we must treat the spatial and Euclidean time extents separately in our analyticity arguments. This follows essentially as a direct consequence of the noncommutativity of the operator elements of the Hamiltonian in the quantum case.

In the quasi-classical series expansion formalism we would now have difficulty proceeding with our discussion as a basis for practical calculations because the basis operations in the extended phase are nonlocal. This means that we could only recover both of our limits simultaneously : the discrete statistical distribution being built up step-by-step is that for the entire volume, rather than for the elementary plaquettes alone. Conversely, the factored product elements we will now focus on are being defined towards the limit of vanishing Euclidean time lattice spacing, thus giving an analyticity ahead of the continuum limit for the spatial extent through a preserved local symmetry. It is this effective treatment of the disconnected contributions that allows us to proceed in the quantum case with the factorised product approach.

What we want to now consider, therefore, is if the eigenvalue spectra, assumed now complete and covering for the extended space, has a meaningful connection with the poles of the free energy density of the partition function and if these poles then demonstrate the lattice volume scaling properties associated with partition function zeroes analysis of Lee, Yang and Fisher. We do this now by noting recent results on the structure of the inverse Laplace transform poles of the transfer matrix elements in the context of discussions on exact probabilistic lattice dynamics. To follow this recent discussion, and deduce our zeroes continuum limit behaviour, we can rewrite the loop operator expression determined for the partition function through the notation of [5][6]. In this case multiplying by the infinitesimal Poissonian probability elements in the Euclidean time

intervals $[t_1, t_1 + dt_1], \dots, [t_N, t_N + dt_N]$ the off-diagonal operator matrix elements $\mathcal{W}_N(\beta)$ of h_b are now expressed as,

$$\mathcal{W}_N(\beta; V_0, V_1, \dots, V_N) = \epsilon^N \left\{ \prod_{i=1}^N \int_0^\beta dt_i \theta(t_i - t_{i-1}) \exp \Delta_i t_i \right\} e^{-V_N \beta} \quad (11)$$

with $\Delta_i = V_i - V_{i-1}$ the potential energy difference between the sites at i and $i-1$, and ϵ some arbitrary energy scale for the purposes of normalisation. In the previous section we showed that in the limit of vanishing Euclidean time lattice spacing we recover a time-ordered Poisson distribution for the elements of the factored spatial basis. Here this is the initial assumption for the product : an element remains in a given basis state until it undergoes a Poissonian distributed event. We can thus consider this the generic form for the loop operator elements in this limit. Here we are again expanding the product over the basis of states N , the extent of which corresponds to the spatial volume, with the integrand here corresponding to the time-ordered limit of the graph distributions. In the discrete case the graph paths can propagate throughout the configuration volume, and equally here the product elements are in causal connection. The above element maps between consecutive spatial sites i , with all four corners of the elementary plaquette considered by definition, and again S is locally conserved. The explicit difference, however, between this weight form $\mathcal{W}_N(\beta)$ and h_b lies in the expression of the specific matrix element contributions. The Heaviside step function being unidirectional over the product indexing here implies that the mapping of $\mathcal{W}_N(\beta)$ defines only one element of the transfer matrix. The time-ordering is here implicit to the potential energy values recovered from the graph determination process, whereas the ordering of the product indexing simply labels the element within the elementary plaquette. Since this latter ordering maps between successive site indices i we can associate $\mathcal{W}_N(\beta)$ with the off-diagonal contribution to the loop operator h_b . Why we believe it relevant to consider just this subset of h_b is that for the generic form of h_b for a quantum system, as we have argued (somewhat tacitly) in the previous section, the diagonal elements form the identity operator. The zeroes of the operator product are thus characterised by the values of the self-adjoint off-diagonal contributions.

Notice again that the above expression is a result for an exactly ergodic Markov process, and so we must have achieved this statistical limit of vanishing Euclidean time lattice spacing for the above relation to be valid. We have argued, somewhat heuristically, that this limit is reached by generating a sufficient number of the formally defined graphs such that these graphs then recover a suitably complete basis for the extended space. Since the transfer matrix elements are as well defined over the ensemble rather than for a single lattice configuration, similarly we will be interested in the zeroes of the ensemble-averaged characteristic polynomial rather than those of a single

configuration for a finite Euclidean time extent. The above relation for the weight form $\mathcal{W}_N(\beta)$ is of course given in terms of potential energies rather than the polynomial zeroes of the loop operator product, but the analyticity of the zeroes and poles are in fact intimately connected through the properties of the inverse Laplace transform of $\mathcal{W}_N(\beta)$, which we will now consider following [5][6]. Differentiating with respect to β and substituting for the Laplace transform $\widetilde{\mathcal{W}}_N(z)$ of $\mathcal{W}_N(\beta)$ reduces the weight form to the following recursive equation,

$$\widetilde{\mathcal{W}}_N(\beta) = \int_0^\infty d\beta e^{-z\beta} \mathcal{W}_N(\beta), \quad z \in \mathbb{C} \quad (12)$$

$$\partial_\beta \mathcal{W}_N(\beta) = \epsilon \mathcal{W}_{N-1}(\beta) - V_N \mathcal{W}_N(\beta), \quad (13)$$

$$\widetilde{\mathcal{W}}_N(z) = \epsilon(z + V_N)^{-1} \widetilde{\mathcal{W}}_{N-1}(z), \quad (14)$$

$$= \epsilon^N \prod_{i=0}^N \frac{1}{z + V_i}. \quad (15)$$

The motivation for introducing these new relations is essentially to quantify the asymptotic scaling behaviour of the expansion via discussion of saddle point solutions. The above poles of the Laplace transform are by definition confined to be in the real segment $[-V_{\max}, -V_{\min}]$, with $V_{\max}$ and $V_{\min}$ respectively the maximum and minimum elements of the set of potential energies $\{V_i\}$. It follows from the definition of the Laplace inverse transformation that its integration contour Γ is any line parallel to the imaginary axis contained in the analyticity domain of the Laplace transform, ie. $\text{Re } z > -V_{\min}$. Applying a saddle point approach to the expansion it can be shown that the first $N - 1$ solutions lie in the range $-V_{\max} < \text{Re } z_0 < -V_{\min}$ with the last solution lying such that $\text{Re } z_0 > -V_{\min}$, where z_0 are the locations of the saddle points and the ordering of solutions is in terms of increasing size. The restrictions on the poles come from quite general considerations, the only real condition here being that β is necessarily positive [20][21].

The whole point or purpose of the saddle point analysis here is that its solution is independent of the path distributions of the graphs but rather dependent on the local properties of the graphs as given through the potential energy differences. From this consideration then the integration contour can be deformed to a new parallel one, although we must know by definition that the path mapped by the weight is analytic along the contour Γ and where the isolated singularities exist. One could therefore then in principle equally well treat the general case of complex-valued potential energy differences, which would be similarly defined since Γ would have no explicit singularities from the local properties of the graphs by construction. Generally, of course, there is no reason to suppose that the poles have to be uniquely defined, and could be given only up to a branch cut, but this is not a problem with the asymptotic form since in the asymptotic limit this choice is naturally resolved. We

should note that through the above definitions and those of the previous section we are at the same time uniquely establishing the residue properties of the weight function and thus the existence of the zeroes of the loop operator factored polynomial.

The last saddle point solution alone is compatible with the restrictions on the integration contour Γ , and can be further shown to be bounded from below by $\text{Re } z_0 > -V_{\min} + N_{V_{\min}}/\beta$ [5]. Thus for $\beta \rightarrow \infty$ with $N_{V_{\min}} \sim N \sim \beta$ the distance between the saddle point and the closest pole remains finite, and the zero temperature continuum limit is well-defined. The distribution of the properly defined complete zeroes now necessarily have some limiting distribution with relevance to the continuum from the asymptotic behaviour of these poles. To be clear, the partition function is expanded in whatever constant diagonal prefactor is introduced in h_b . It is proportional to β for the pedagogical $S = 1/2$ AFM example and therefore the analysis is specifically of interest for (although not restricted to) the case of quantum fluctuations at low temperatures. Strictly, as we have seen, only one of the poles lies in the analyticity domain of the polynomial. This does not imply that the characteristic polynomial is undefined, but rather that we only understand the asymptotic behaviour of the weight function in the vicinity of this one pole. There is a circle theorem by Lee and Yang which we will now discuss which relates to the limiting behaviour of branch cuts, which we can perhaps envisage now being applicable in at least one instance of the zero temperature limit. To reiterate, the noncommuting nature of the quantum system Hamiltonian has forced us to consider a limiting Euclidean time spacing distribution defined over the spatial basis of our lattice, for which at least one of the zeroes of the polynomial by construction has a well defined continuum limit.

IV. SCALING OF PARTITION FUNCTION ZEROES

We have considered how a partition function can be defined over an extended phase space for a quantum system through a loop operator formalism, and also how the limit of vanishing Euclidean time lattice spacing is related to the analyticity properties of the characteristic polynomial of this factored product and its pole structure. Although the recurrence relations defined for the above Laplace transforms of the operator elements have an explicit dependence on the spatial lattice volume N , we must further explore this connection to firmly establish the zeroes content regarding the singularities of the partition function as the limit $N \rightarrow \infty$ is approached. The starting point of Lee and Yang's arguments is essentially that a valid zeroes polynomial form exists, from which finite volume scaling discussion can be then made [2]. We have now argued that the analyticity of factorisation property of a quantum partition function is in general not possible without first approaching an asymptotic scaling

regime : a far weaker precondition for this zeroes form. Comparing with the constraints we have thus far defined on the factored product eigenvalues for the quantum system we now contrast with the two central criterion of Lee and Yang's discussion for defining valid thermodynamic functions from the partition function in the limit $N \rightarrow \infty$ of the spatial lattice volume [2].

The first of these criterion is that the limiting value of as the pressure of the system P , is independent of the shape of N and is a continuous monotonically increasing function of the expansion parameter of the suitably factored partition function.

$$P = \lim_{N \rightarrow \infty} \frac{1}{N} \log(Z) \quad (16)$$

Since the action of the factorisation in the quantum case is by definition local and ergodic it follows that the weights are obtained recursively, as we have seen. The limit $(N+M) \rightarrow \infty$ of the difference of the log of two consecutive intensive weights of integer index N and $(N+M)$ is thus zero. Establishing this property one can then consider the volume as consisting of a finite subset of smaller volume elements, with interactions across the boundaries between subsets for which the free energy of this interface tension does not grow faster than the surface area. The above result thus implies that the limit of thermodynamic functions, including the pressure, is always finite. Monotonicity and the continuous nature of the functions follows from the similar behaviour of the derivatives of the partition function, as is seen explicitly through the properties of the inverse Laplace transform. Thus the distribution of the zeroes necessarily gives all the analytic behaviour of thermodynamic functions in the expansion parameter plane as $N \rightarrow \infty$.

The second of Lee and Yang's criterion is that the limiting value of the density of the system ρ , is an increasing function of the expansion parameter of the suitably factored partition function but does not necessarily take a defined limit for all expansion parameter values, β .

$$\rho = \lim_{N \rightarrow \infty} \frac{\partial}{\partial \log(\beta)} \frac{1}{N} \log(Z) \quad (17)$$

Again this is very much an implicit result for the factored product of the quantum system Laplace transform and the discussions of the previous section . The criterion implies that if a region of the positive real axis of β is free of zeroes of the partition function as $N \rightarrow \infty$ then the system is here in a single phase. The proof of this follows from noticing that the coefficients of the factored product expansion are necessarily positive, which is a fairly general consequence of the operators being self-adjoint in the quantum case.

Lee and Yang's first discussion with their new result was essentially to notice that there was necessarily a problem in discussing thermodynamic functions with solely the

$N \rightarrow \infty$ limit of the partition function polynomial expansion coefficients : this leads to the result of undefined phases. It was not possible to discuss thermodynamic singularities directly in the asymptotic limit, rather the convergence to that limit was the important consideration. They reviewed an earlier treatment of Mayer in which they showed it becomes impossible to analytically continue the series expansion thus formed between phases. This is unlike of course, by construction, the above implicit spatial lattice volume dependencies of P and ρ given from the properties of the Laplace transforms of the operator elements that do yield well defined thermodynamic functions. The difference lies in the differing properties of a series and polynomial. The series expansion values do not yield a density dependence to pressure for specific volumes below the point at which the zeroes approach the real β axis (the phase transition point). This is implicit since the continuum values have obviously no spatial lattice volume dependence. One can therefore argue from this that not only is the partition function zeroes form for the quantum system compatible with Lee and Yang's discussion, but also that it is therefore in some sense a necessary precondition for all partition function zeroes expansions, classical and quantum. To expand : we have laboured to ensure the factored product is meromorphic, singular only at the poles, which are themselves in a one-to-one correspondence with the zeroes. A loop operator picture defined over an extended phase space must be therefore factorisable for all the thermodynamic functions to be able to defined for a system, since it is only the existence of the limit of vanishing Euclidean time lattice spacing that ensures that the polynomial expansion zeroes, and by extension the expansion coefficients, have a valid asymptotic form.

We can generalise this discussion further to consider the implications for complex operator elements and the introduction of symmetry breaking source terms defined with relation to the limit $N \rightarrow \infty$. Partition function zeroes have been recently used to discuss the Vafa-Witten theorem and complex action problems for quantum systems [22]. To summarise this discussion : the Vafa-Witten result is that parity and CT cannot be spontaneously broken in vector-like theories such as QCD. The argument follows from two points ; any arbitrary local hermitian order parameter for parity X , constructed from Bose fields is proportional to an odd power of the totally antisymmetric tensor, and the free energy density is well-defined in the presence of a symmetry breaking source θX , with θ some real constant. These two conditions jointly then imply that the symmetry breaking source term be a phase in the integrand of the Euclidean partition function. The explicit connection of the theorem with partition function zeroes is then made by considering $p(\tilde{X}, N)$ the probability density function of $\tilde{X}$, where $\tilde{X}$ expresses X extensively. Assuming parity to be spontaneously broken $p(\tilde{X}, N)$ can then be argued to be taken as the sum of two delta functions in the limit $N \rightarrow \infty$, centred on $\pm a$. The relevant zeroes of the parti-

tion function in θ are thus obtained as the solutions of,

$$\cot(\theta Na) = \frac{\int_0^\infty d\tilde{X} p(\tilde{X}, N) \sin(\theta N(\tilde{X} - a))}{\int_0^\infty d\tilde{X} p(\tilde{X}, N) \cos(\theta N(\tilde{X} - a))}. \quad (18)$$

This has an infinite number of solutions approaching the origin, and so the Vafa-Witten theorem is thus seen to be upheld since the second tenet of free energy is then undefined. In fact this is a general consequence of the properties of the Laplace transforms of complex-valued integrands. The asymptotic convergence, as a function of θ , is in general not defined for complex-valued functions given for the exponent : the oscillatory phase means the integration contour Γ is ill-defined. Conversely, as we have argued, following Lee and Yang's discussion of Mayer's series treatment it is possible to consider the symmetry breaking consequences of a complex-valued system if the expansion coefficients are expressed intensively. Namely, if we have a contribution to the partition function that is highly nonlocal but preserves an exact local symmetry the asymptotic convergence properties of the partition function are defined, as we have seen explicitly with the above loop operator formalism.

We can understand in a very simple way how singularities of the partition function arise as $N \rightarrow \infty$ from our pedagogical loop operator example. Consider a self adjoint matrix with real elements of values between zero and one and the product of all elements in the upper triangular region such that these give recursively the characteristic polynomial expansion coefficients to successive order [23]. It can be shown by induction that if these elements have this probabilistic interpretation then necessarily the zeroes of the characteristic polynomial are bounded on the unit circle. In the case of Lee and Yang's discussion for the Ising model with external field we can then identify the field associated with generating the off diagonal transfer matrix elements and a symmetry breaking nonzero source value. Although here there is no such association we can still understand the singularities of the spatial extent of the partition function in the same way, and therefore can similarly associate a critical value to our factored product expansion parameter with a valid asymptotic form. Generalising for complex valued operator elements we see that the critical value of coupling is still recovered through a similar boundedness induction argument as the operators are necessarily hermitian in this case. We have in the loop operator introduction discussion and subsequent factorised product discussion essentially defined a source term such that it is only recovered in the asymptotic limit, thus obviating the ambiguity of contour definition for the Laplace transform for the complex case.

Our initial motivations included a comparison of quasi-classical and factored product expansions for the treatment of quantum systems. It is perhaps now more evident why we have favoured the latter for this zeroes

treatment : the complex action problem above being amenable to treatment via quantum fluctuations. The equivalence of the quasi-classical approach to the continuum is only resolved when both spatial and time extent limits are simultaneously reached, but more over the symmetry breaking mechanisms are otherwise potentially contaminated with disconnected contributions to the partition function. Conversely, the nonlocal nature of the intermediate result of Euclidean time lattice spacing for the factored product case means that nonlocal symmetry breaking terms can be considered but on a countably finite space, yielding well-defined thermodynamic functions in the continuum. The quantum fluctuations represented through the partition function are inherently nonlocal and it is only through this latter limiting formalism that the disconnected contributions are exactly articulated since otherwise, as we have seen, the generating functional for the partition function becomes simply a generating function.

There is an intrinsic connection between this discussion and the renormalization group arguments made by Wilson through the implicit differentiability conditions introduced for the spatial lattice volume extent [24]. Looking at the density relation, the condition used to establish the $N \rightarrow \infty$ zeroes singularities, it is possible to understand that the critical points on the real axis of the expansion parameter correspond to the fixed points of the renormalization group equation defined in the limit $N \rightarrow \infty$. This is a new result of the loop operator treatment since for the general classical system partition function zeroes case it is not possible to make this association, as the asymptotic scaling properties of the expansion parameter are not defined. The expansion parameter in this case can be strongly dependent on the Euclidean time lattice spacing and singular and dependent on the asymptotic properties therein. The differentiability conditions of the renormalization group equations have an obvious expression in the recursion relation for the Laplace transforms we have discussed, and the group property can be understood in this mapping between successive poles. Returning to the Vafa-Witten comment it is of course possible as with the renormalization group discussion that discontinuities between minima of the vacua arise in the asymptotic limit, in which case it is possible to discuss phase transitions even in the case that we have no intensive source term comparing with the Ising model with external field [22]. The difference in level between the peaks of the vacua essentially then does not have to be infinite in order that an exact discontinuity arises in the asymptotic limit of the expansion parameter β which would now play an analogous role to time in the motivation of the renormalization group equations.

In this loop operator treatment it is therefore important to understand the evolution of the spatial lattice spacing dependence in regards to the nonlocal contributions to the expansion : infrared divergences potentially lie in the nonlocality of the expansion. It is important to understand as well how the singularities of the expan-

sion relate to symmetry breaking and the emergence of Goldstone modes which we will now discuss.

V. ASYMPTOTIC FREEDOM

The properties of branch cuts are clearly important in the partition function zeroes singularity picture as they relate to the emergence of a form of geometric bound on the modulus and argument of the zeroes in the complex expansion parameter plane [23]. It is only the property of the zero nearest the real axis which is strictly important in determining the singularities in the $N \rightarrow \infty$ limit, but in the quantum case this also establishes the behaviour of an asymptotically defined source for the singularity. The scaling of the feature is therefore critically important to determining whether or not singularities of the zeroes in the spatial lattice polynomial expansion correspond to fixed points of this renormalization group equation defined along the real axis in β . The sense in which we associate the loop operator partition function zeroes treatment to the renormalization group equation approach is perhaps therefore closer to discussions of the strong interaction and a momenta dependent treatment [4]. Although the spatial volume is clearly important we are not treating the lattice spacing dependencies of this extent in a direct manner, and so the $N \rightarrow \infty$ properties of the zeroes are in a sense disconnected from the asymptotic treatment of the expansion parameter : in the sense that we would be interested to relate the correlation length and bare parameters of the system to finite N scaling properties of the volume.

Never the less the symmetry breaking mechanisms of the system can be understood tentatively with relation to the Euclidean time extent and correlation length. If we take a conventional renormalization group analysis of a system like the 2-d $O(3)$ model we can understand that the correlation length, relating to the conventional renormalization group β -function coefficients, diverges exponentially in the limit of vanishing lattice spacing and so there are massless particles [1]. However, for this particular system, the Mermin-Wagner-Coleman theorem prevents the massless Goldstone bosons interacting and so the model has a nonperturbatively generated mass gap [25]. Conversely other systems can be pictured in which the correlation length remains infinite and we have a broken symmetry, but reducing the Euclidean time extent β eventually recovers a symmetric phase with a finite correlation length. If the Euclidean time extent is very much smaller than the correlation length we can consider that all these systems undergo dimensional reduction : we can treat one direction as effectively static. This then allows the continuum limit of these models to be defined with $\beta \rightarrow \infty$. This limit is of interest because we are interested in the asymptotic properties of β : which fixed point, or singularity on the real axis, we end up at in this limit. Notice that we have not aimed to fix at any point an association of the loop operator

formalism with this dimensional reduction. In essence we therefore invert the above argument : having determined singularities in the limit $N \rightarrow \infty$ we then consider the properties of these points on the real β axis where asymptotic convergence to the continuum is in principle guaranteed by construction. This is particularly useful form for the direct analysis of numerical lattice results since it is possible to understand from direct examination of the singularities whether or not they correspond to genuine fixed points of a given model or are simply lattice spacing dependent cutoff effects.

Considering the arguments defining a renormalization program for QED to all orders perturbatively in terms of momenta, one must at some point make momentum cutoff dependent subtractions in the diagrams [4]. Prescriptively constants are chosen so that the relevant Ward identities are satisfied at one particular value of momentum, and such that the renormalization functions do not contain singularities for the zero electron mass case. Similarly by construction the renormalized amplitudes of the diagrams do not contain infrared divergences for zero photon mass and nonzero electron mass. This prescription for infrared divergences of Gell-mann and Low's discussed corresponds to the asymptotic limit of momentum cutoff being much bigger than the mass, and the solutions of the renormalization group equations for all momentum cutoff values and zero photon mass necessarily then define a scale invariant field theory. One can break this scale invariance in one of two ways ; by letting the mass be nonzero or by choosing a solution of the renormalization group equations at zero mass that is not a fixed point. One can therefore understand by analogy that for the loop operator approach all singularities in the spatial N extent can be connected to a finite β value with an asymptotic form, but that these points don't necessarily correspond to fixed point solutions if they do not have any support. To elucidate slightly : if a zero has an edge singularity associated with boundedness then its form is akin to this former case and we would have a broken symmetry with a correlation length that diverges as the continuum limit is approached, and the converse is that of a finite correlation length with a valid continuum limit. The third case is that of the mass gap, which would be otherwise under-defined in the analysis of say the case for example of the Ising model with external field, since some points of the complex phase plane diagram are necessarily undefined with the Laplace transform of the poles having essentially an under-defined contour [22].

Consider as an example the Haldane conjectured antiferromagnetic $O(3)$ quantum spin chains, where integer spins have a mass gap and half-integer spins are gapless [1]. The low-energy effective theory for this system is the 2-d $O(3)$ model at vacuum angle $\theta = \pi$, which is in the universality class of the $k = 1$ Wess-Zumino-Witten model. Numerical studies of this model indicate that the gap which is present for all $\theta \neq \pi$, vanishes at the specific value $\theta = \pi$. When the Euclidean time extent β is made finite this topological term also disappears. One can con-

clude from this system therefore that one would expect to see a singularity on the β axis from the loop operator formalism, but that this should not scale directly to the continuum because the finite volume dependence of the scheme essentially introduces a divergent correlation length and one sees otherwise a pseudo-critical scaling. However the lattice singularity still exists on the finite volume and its relation to the asymptotic scaling can be measured directly from the lattice result, since again one can obviously study the gap dependence on finite system size directly in numerical simulation. The advantage with the new loop operator method is firstly that we can quantify the directly the asymptotic scaling properties of the gap itself, but also understand the evolution of other cutoff dependent quantum phase transition effects from the scaling arguments of related singularities.

Preliminary results for the loop operator zeroes scheme are presented for the case of mixed integer - half-integer spin chains which share a related gapless property to the integer chains but with an additional saddle point effect [26]. These admit a natural probabilistic loop operator treatment as well following the pedagogical $S=1/2$ AFM treatment schemes we have followed and discussed. An exact treatment has been recently given for the dimensional reduction aspects of plaquette defined operators over an extended phase space for the case of gauge field theories. This raises then raises the possibility of performing a similar analysis to the loop operator treatment for the case of gauge fields, as opposed to quasi-classical treatment, but the issues of numerically simulation are by no means trivial to address. A general pathology of the factorised loop-cluster method, where the Monte Carlo decisions are taken plaquette wise, is the statistical limitations of building up a long tailed probability distribution, and extending to the case of gauge or complex valued fields one could then have also non positive definite operator elements which do not yield a probabilistic interpretation at all. One option is to use a projection operator basis acting on a conventional globally updated space to define pseudo-plaquette states [27][28]. Strictly from the Laplace transform arguments we have reviewed and presented the only necessary criterion for a plaquette term on the space is that it contain at most one pole and have an continuous probability density function. In QCD, for example, a basis of related operators can be compiled over a given lattice gauge field ensemble such that the poles of the basis suppress unphysical state contributions through a natural ranking [29]. Here we define first a Euclidean time dependent polynomial form for the global update scheme, independent of the related operator basis, and introduce a limiting product such that the elements of these pseudo-plaquette operators contain at most one zero of the polynomial, which is easily done since the probability density functions of the operators on the globally generated gauge ensemble become universally distributed at some critical scale [30]. One can understand this latter result from general consideration of the central limit theorem in regard to defining the con-

tour for the above Laplace transforms, and its relation to the stationary phase approximation in this regard [31].

In lattice field theory calculations one can avoid some of the uncertainties associated with truncating a perturbative expansion in determining constants in the renormalization group equations by defining various nonperturbative schemes [32][33]. Constants are set through the chiral Ward identities for the lattice Green functions. For example with RI/MOM the conditions are set on Green functions at large external momenta, while in the Schrödinger Functional scheme they are set in small lattice volumes. The lattice renormalization scales have to be larger than Λ_{QCD} , to connect with perturbation theory and smaller than the inverse lattice spacing to avoid cutoff effects, which is perhaps computationally expensive. With the RI/MOM approach as well the pseudoscalar Green function, which couples to the Goldstone boson, has a leading contribution diverging in the chiral limit. Similarly in building candidate supersymmetric lattice models one is interested in the Ward identities in the continuum limit in terms of renormalized perturbation theory [34]. Schemes involve, to finite order in perturbation theory, adjusting a coefficient in local terms in the action but here as well it is possible for the action to become complex [35]. In both cases it is possible to envisage calculating a simple limiting operator factorisation with some Euclidean time extent polynomial scheme defined such that the asymptotic properties of these lattice renormalization properties could be further studied directly in this context.

VI. SUMMARY

We have discussed, in the context of recent ideas on the rôle of dimensional reduction in quantum systems and reviews of loop operator methods a formulation of a partition function polynomial zeroes treatment for quantum systems. We have argued that in general the quantum case should be treated separately from the classical case through a local projection operator argument for the factored product, suggesting the evolution of infrared divergences is otherwise implicit to the quantum case. We have relaxed the condition that the expansion variable be expressed through an exponential function to this end, whilst still retaining its asymptotic structure. By working over an extended phase space we have then identified a limiting form with a connection to the continuum, and the fixed point solutions of renormalisation group equations, using recent results for the treatment of exact probabilistic lattice dynamics. Related to this continuum limiting process we have then discussed the prospects for further evaluating gauge field theories for the purposes of perturbation theory independent renormalization group analysis, and the treatment of theories with a non positive-definite source terms in the Euclidean action. The outline is necessarily general but further investigation of the limiting analyticity of the partition

function is proposed, with three cases with relevance to symmetry breaking mechanisms envisaged as pertinent

for future analysis and categorisation.

-
- [1] R. Brower, S. Chandrasekharan, S. Riederer and U.-J. Wiese, Nucl. Phys. B **693** (2004) 149.
 - [2] T.D. Lee and C.N. Yang, Phys. Rev. **87** (1952) 404.
 - [3] D.A. Kurtze and M.E. Fisher, Phys. Rev. B **20** (1979) 2785.
 - [4] K.G. Wilson, Phys. Rev. D **3** (1971) 1818.
 - [5] M. Ostili and C. Presilla, [[hep-lat/0412157](#)].
 - [6] M. Ostili and C. Presilla, New J. Phys. **6** (2004) 107.
 - [7] H.G. Evertz, Adv. Phys. **52** (2003) 1.
 - [8] S. Todo and K. Kato, Phys. Rev. Lett. **87** (2001) 047203.
 - [9] M. Troyer, F. Alet, S. Trebst and S. Wessel, AIP Conf. Proc. **690** (2003) 156.
 - [10] B.B. Beard and U.J. Wiese, Nucl. Phys. Proc. Suppl. **53** (1997) 838.
 - [11] A.W. Sandvik, Phys. Rev. B **59** (1999) R14157.
 - [12] S. Gottlieb, W. Liu, D. Toussaint, R.L. Renken and R.L. Sugar, Phys. Rev. D **35** (1987) 2531.
 - [13] K. Kladko, P. Fulde and D.A. Garanin, Europhys. Lett. **46** (1999) 425.
 - [14] P. Bialas, A. Morel, B. Petersson, K. Petrov and T. Reisz, Nucl. Phys. B **581** (2000) 477.
 - [15] S. Kratochvila and P. de Forcrand, Nucl. Phys. Proc. Suppl. **106** (2002) 522.
 - [16] A. Cuccoli, T. Roscilde, V. Tognetti, R. Vaia and P. Verucchi, Phys. Rev. B **67** (2003) 104414.
 - [17] R.H. Swendsen and J.S. Wang, Phys. Rev. Lett. **58** (1987) 86.
 - [18] M. Aizenman, B. Nachtergaele, Commun. Math. Phys. **164** (1994) 17.
 - [19] Y. Nambu, Phys. Rev. D **7** (1973) 2405.
 - [20] J. Kevorkian and J.D. Cole, *Multiple Scale and Singular Perturbation Methods*, Volume 114 in the Applied Mathematical Sciences Series, Springer Verlag, Berlin, 1996.
 - [21] J.D. Murray, *Asymptotic Analysis* Volume 48 in the Applied Mathematical Sciences Series, Springer Verlag, Berlin, 1984.
 - [22] V. Azcoiti, V. Laliena and A. Galante, in *Guangzhou 2000, Nonperturbative methods and lattice QCD*, edited by X.Q. Luo and E.B. Gregory (Singapore, World Scientific, 2001). [[hep-lat/0007045](#)].
 - [23] T.D. Lee and C.N. Yang, Phys. Rev. **87** (1952) 410.
 - [24] K.G. Wilson, Phys. Rev. B **4** (1971) 3174.
 - [25] C. Itoi, [[hep-lat/0303118](#)].
 - [26] P.R. Crompton, W. Janke, Z.X. Xu and H.P. Ying, Nucl. Phys. Proc. Suppl. **140** (2005) 817.
 - [27] P.R. Crompton, Nucl. Phys. B **619** (2001) 499.
 - [28] P.R. Crompton, Nucl. Phys. B **626** (2002) 228.
 - [29] D. Brömmel, P.R. Crompton, C. Gattringer, L.Y. Glozman, C.B. Lang, A. Schäfer and S. Schäfer, Phys. Rev. D **69** (2004) 094513.
 - [30] P.R. Crompton, Nucl. Phys. Proc. Suppl. **119** (2003) 550.
 - [31] P.R. Crompton, in *Strong and Electroweak Matter 2002: Proceedings*, edited by M.G. Schmidt (River Edge, N.J., World Scientific, 2003). [[hep-lat/0301001](#)].
 - [32] D. Bécirévic, V. Gimenez, V. Lubicz, G. Martinelli, M. Papinutto and J. Reyes, JHEP **0408** (2004) 022.
 - [33] V. Gimenez, L. Giusti, S. Guerriero, V. Lubicz, G. Martinelli, S. Petrarca, J. Reyes, B. Taglienti and E. Trevigne, Phys. Lett. B **598** (2004) 227.
 - [34] S. Catterall, JHEP **0411** (2004) 006.
 - [35] Y. Kikukawa and H. Suzuki, JHEP **0502** (2005) 012.